

## Erratum to: Evaluation of physical activity and its relationship to health-related quality of life in patients on chronic hemodialysis

Akihiko Katayama · Nobuyuki Miyatake ·  
Hiroyuki Nishi · Kazuhiro Uzike · Noriko Sakano ·  
Hiroo Hashimoto · Kiichi Koumoto

Published online: 7 February 2015  
© The Japanese Society for Hygiene 2015

**Erratum to: Environ Health Prev Med (2014)  
19:220–225  
DOI 10.1007/s12199-014-0380-z**

The authors would like to correct the error occurred in the ‘Result’ section, of the original publication as below:

Among all patients on chronic hemodialysis, only physical activity over 4 METs on non-hemodialysis treatment days was a determinant factor of EQ-5D [EQ-5D scores = 0.300 (physical activity over 4 METs on non-hemodialysis treatment days) + 0.695,  $r^2 = 0.181$ ,  $p = 0.0026$ ].

---

The online version of the original article can be found under doi:[10.1007/s12199-014-0380-z](https://doi.org/10.1007/s12199-014-0380-z).

---

A. Katayama (✉) · N. Miyatake · N. Sakano  
Department of Hygiene, Faculty of Medicine, Kagawa  
University, 1750-1, Miki, Kagawa 761-0793, Japan  
e-mail: [kata@med.kagawa-u.ac.jp](mailto:kata@med.kagawa-u.ac.jp)

H. Nishi · K. Uzike · H. Hashimoto · K. Koumoto  
Innoshima General Hospital, Onomichi, Hiroshima 722-2323,  
Japan